

Series Representation of the Modified Bessel Functions*

Krzysztof Maślanka

Astronomical Observatory of the Jagiellonian University
Orla 171, 30-244 Kraków, Poland

April 7, 1998

Abstract

Some power series representations of the modified Bessel functions (McDonald functions K_α) are derived using the little known formalism of fractional derivatives. The resulting summation formulae are believed to be new.

1 Fractional derivatives

There are several non-trivial examples in mathematics when some quantity, originally defined as integer, can radically extend its original range and assume fractional or even continuous values. The most common example is the gamma-function of Euler which is a natural, unique generalization of the ordinary factorial:

$$n! \equiv \prod_{i=1}^n i = \Gamma(n+1) = \int_0^\infty t^n e^{-t} dt \quad (n > -1)$$

The same thing may be performed with the order of derivatives which can also be made fractional. Although useful, fractional derivatives do not however create any essentially new calculus being rather some sort of particular, relatively simple, integral transforms.

*Cracow Observatory preprint, no. 7/98

Following Oldham and Spanier (see e.g. [1]; cf. also [2], [3] and [4]) we define the fractional derivative $\partial_{x-a}^s \equiv \left(\frac{\partial}{\partial x}\right)^s$ by an integral representation known as the Riemann-Liouville integral. Given a real number $s < 0$, define

$$\partial_{x-a}^s f(x) = \frac{1}{\Gamma(-s)} \int_a^x (x-t)^{-s-1} f(t) dt \quad (1)$$

where $a < x$ is a fixed number, referred to as the boundary point. Since $s < 0$, the integral is convergent, provided that f behaves well. For $s \geq 0$ we define

$$\partial_{x-a}^s \equiv \partial_x^n \partial_{x-a}^{s-n}$$

where n is a positive integer chosen large enough so that $s-n < 0$ in order to assure convergence of the integral in definition (1). It is not difficult to show that, as expected, such a definition does not depend on n . One can further prove that the familiar Leibniz rule for product differentiation has the form

$$\partial_{x-a}^s (fg) = \sum_{j=0}^{\infty} \binom{s}{j} \partial_{x-a}^{s-j} f \partial_{x-a}^j g \quad (2)$$

where the generalized binomial is

$$\binom{s}{j} \equiv \frac{\Gamma(1+s)}{j! \Gamma(1+s-j)} = \frac{(-1)^j \Gamma(j-s)}{j! \Gamma(-s)}$$

It can also be shown that the following rule for the arbitrary order derivative of the power function holds

$$\partial_{x-a}^s (x-a)^p = \frac{\Gamma(p+1)}{\Gamma(p+1-s)} (x-a)^{p-s} \xrightarrow{s=n} \frac{p!}{(p-n)!} (x-a)^{p-n} \quad (3)$$

for any s and $p > -1$. However, the analogous formulae in the case of the exponential and logarithm functions may, on the first sight, be quite surprising

$$\frac{\partial^s}{\partial x^s} \exp(\beta x) = \beta^s \exp(\beta x) \frac{\gamma(-s, \beta x)}{\Gamma(-s)} \xrightarrow{s=n} \beta^n \exp(\beta x)$$

$$\frac{\partial^s}{\partial x^s} \ln x = \frac{x^{-s}}{\Gamma(1-s)} \left[\ln x - \psi(-s) - C + \frac{1}{s} \right] \xrightarrow{s=n} \frac{(-1)^{n-1} (n-1)!}{x^n}$$

(Here n denotes a positive integer. For brevity we have set the boundary point $a = 0$.)

2 Some definite integrals as fractional derivatives

Let us now consider the following definite integrals (cf. [5], formulae 3.471.4 and 3.471.8)

$$\int_0^x t^{-2\mu} (x-t)^{\mu-1} \exp\left(-\frac{\beta}{t}\right) dt = \frac{\beta^{\frac{1}{2}-\mu}}{\sqrt{\pi x}} \exp\left(-\frac{\beta}{2x}\right) \Gamma(\mu) K_{\mu-\frac{1}{2}}\left(\frac{\beta}{2x}\right) \quad (4)$$

$$\int_0^x t^{-2\mu} (x^2-t^2)^{\mu-1} \exp\left(-\frac{\beta}{t}\right) dt = \frac{1}{\sqrt{\pi}} \left(\frac{2}{\beta}\right)^{\mu-\frac{1}{2}} x^{\mu-\frac{3}{2}} \Gamma(\mu) K_{\mu-\frac{1}{2}}\left(\frac{\beta}{x}\right) \quad (5)$$

which are valid for $x > 0$, $\operatorname{Re} \beta > 0$, $\operatorname{Re} \mu > 0$. On using (1) it is evident that both (4) and (5) may simply be interpreted as fractional derivatives of the appropriate functions. Indeed

$$\frac{\partial^s}{\partial x^s} \left[x^{2s} \exp\left(-\frac{\beta}{x}\right) \right] = \frac{\beta^{s+\frac{1}{2}}}{\sqrt{\pi x}} \exp\left(-\frac{\beta}{2x}\right) K_{s+\frac{1}{2}}\left(\frac{\beta}{2x}\right) \quad (6)$$

$$\frac{\partial^s}{\partial x^s} \left[x^{s-\frac{1}{2}} \exp\left(-\frac{\beta}{\sqrt{x}}\right) \right] = \frac{2}{\sqrt{\pi}} \left(\frac{\beta}{2}\right)^{s+\frac{1}{2}} x^{\frac{3}{4}-\frac{s}{2}} K_{s+\frac{1}{2}}\left(\frac{\beta}{x}\right) \quad (7)$$

It is thus possible even to *define* the functions K_ν as fractional derivatives with zero boundary point.

3 Polynomials V_k

In order to evaluate the left hand side of (6) and (7) let us introduce

$$V_k^{(\alpha)}(\beta x^\alpha) \equiv x^k \exp(\beta x^\alpha) \frac{\partial^k}{\partial x^k} \exp(-\beta x^\alpha) \quad (8)$$

It is easy to prove that, contrary to their appearance, V_k are just polynomials

$$V_k^{(\alpha)}(z) = \sum_{j=0}^k A_{kj}^{(\alpha)} z^j$$

with coefficients given by

$$A_{kj}^{(\alpha)} = (-1)^k \sum_{i=0}^j \frac{(-1)^i}{i! (j-i)!} \frac{\Gamma(k-\alpha i)}{\Gamma(-\alpha i)}$$

In particular, for $\alpha = -1$ we have

$$A_{kj}^{(-1)} = (-1)^k \sum_{i=0}^j \frac{(-1)^i}{i! (j-i)!} \frac{\Gamma(k+i)}{\Gamma(i)} = \frac{(-1)^{k+j} k! (k-1)!}{(k-j)! j! (j-1)!}$$

(The last equality may be rigorously proved using elementary methods presented in e.g. [7]; *Mathematica* 3.0 effectively simplifies such sums.) We shall also need fractional derivatives of the following expressions

$$x^\nu \exp(-\beta x^\alpha)$$

with α, ν real. Using Leibniz rule (2) and the property (3) we have

$$\frac{\partial^s}{\partial x^s} [x^\nu \exp(-\beta x^\alpha)] = x^{\nu-s} \frac{\Gamma(\nu+1)}{\Gamma(-s)} \exp(-\beta x^\alpha) \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \frac{\Gamma(k-s)}{\Gamma(k-s+\nu+1)} V_k^{(\alpha)} \quad (9)$$

4 Series expansion of the Modified Bessel functions

Inserting (9) into (6) or (7) we can get a general expansion of the modified Bessel functions (called also McDonald functions; cf. e.g. [6]) holding for any positive α

$$K_s(z) = \frac{\sqrt{\pi}}{(2z)^s} \exp(-z) \frac{\Gamma(2s)}{\Gamma(\frac{1}{2}-s)} \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \frac{\Gamma(k+\frac{1}{2}-s)}{\Gamma(k+\frac{1}{2}+s)} V_k^{(-1)}(2z) \quad (10)$$

Substituting polynomials (8) into (10) we further get, after some manipulations

$$K_s(z) = 2^{s-1} \Gamma(s) z^{-s} \exp(-z) \left[1 + \sum_{k=1}^{\infty} \frac{\left(\frac{1}{2}-s\right)_k}{\left(\frac{1}{2}+s\right)_k} \sum_{j=1}^k \binom{k-1}{j-1} \frac{(-2z)^j}{j!} \right]$$

where

$$(a)_k \equiv \frac{\Gamma(k+a)}{\Gamma(a)}$$

denotes the Pochhammer symbol. The same method applied to (7) gives

$$K_s(z) = 2^{s-1} \sqrt{\pi} \frac{\Gamma(2s)}{\Gamma(\frac{1}{2}-s)} z^{-s} \exp(-z) \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \frac{\Gamma(k+\frac{1}{2}-s)}{\Gamma(k+\frac{1}{2}+s)} V_k^{(-1/2)}(z) \quad (11)$$

which is yet another power series convergent to the McDonald function. The familiar ratio of Pochhammer symbols in (10) and (11) allows us to call these expansions 'hypergeometric-like' since they formally resemble the well-known hypergeometric Kummer function ${}_1F_1$.

References

- [1] K. B. Oldham, J. Spanier, *The Fractional Calculus*, Academic Press, New York, 1974
- [2] V. S. Vladimiroff, *Obobshchennye funktsii v matematicheskoy fizike*, Nauka, Moskva, 1979 (in Russian)
- [3] R. Camporesi, *Harmonic Analysis and Propagators on Homogeneous Spaces*, Physics Reports 196 no. 1&2, 1990
- [4] K. Maślanka, *Efekty kwantowe w wielowymiarowych modelach kosmologicznych*, Ph. D. thesis, 1986, (unpublished, in Polish)
- [5] I. S. Ryzhik, I. M. Gradstein, *Tablicy intiegralov, sum, riadow i proizvedenij*, Nauka, Moskva, 1971, (in Russian)
- [6] M. Abramowitz, I. A. Stegun, eds., *Handbook of Mathematical Functions*, U.S. National Bureau of Standards, Dover, New York, 1965
- [7] R. L. Graham, D. E. Knuth, O. Patashnik, *Concrete Mathematics. A Foundation for Computer Science*, Addison-Wesley, 1994